Brain Research, 462 (1988) 407–409
Elsevier

Author Index

Abraham, W.C. and Mason, S.E., Effects of the NMDA receptor/channel antagonists CPP and MK801 on hippocampal field potentials and long-term potentiation in anesthetized rats, 40
Åkerman, K.E.O., see Enkvist, M.O.K., 67
Alkon, D.L., see Disterhoft, J.F., 118
Alvarez, F.J., see Villalba, R.M., 359
Amir, S. and Jackson, I.M.D., Immunological blockade of endogenous thyrotropin-releasing hormone impairs recovery from hyperglycemia in mice, 160
Arrezo, J.C., Brosnan, C.F., Schroeder, C.E., Litwak, M.S. and Bornstein, M.B., Electrophysiological analysis of factors involved in the primary demyelinating diseases: the rabbit eye model system, 286
Aschner, M. and Clarkson, T.W., Uptake of methylmercury in the rat brain: effects of amino acids, 31

Baker, M., see Bostock, H., 354
Becker, J.B. and Freed, W.J., Adrenal medulla grafts enhance functional activity of the striatal dopamine system following substantia nigra lesions, 401
Belluardo, N., see Bindoni, M., 10
Bennett, J.A., see Robinson, T.E., 211
Bentgen, K.M., see Robinson, T.E., 211
Bernardi, G., see Mercuri, N.B., 199
Berretta, S., see Bindoni, M., 10
Bickford-Wimer, P.C., see Parfitt, K.D., 242
Bicknell, R.J., see Zhao, B.-G., 62
Bindoni, M., Perciavalle, V., Berretta, S., Belluardo, N. and Diamantstein, T., Interleukin 2 modifies the bioelectric activity of some neurosecretory nuclei in the rat hypothalamus, 10
Blasco, I., see Villalba, R.M., 359
Borgers, M., see Van Reempts, J., 142
Bornstein, M.B., see Arrezo, J.C., 286
Bortollotto, Z.A., see Ikonomidou-Turski, C., 275
Bostock, H. and Baker, M., Evidence for two types of potassium channel in human motor axons in vivo, 354
Brosnan, C.F., see Arrezo, J.C., 286

Calabresi, P., see Mercuri, N.B., 199
Calderazzo-Filho, L.S., see Ikonomidou-Turski, C., 275
Calvet, J., see Calvet, M.-C., 321
Calvet, J., see Calvet, M.-C., 334
Calvet, M.-C. and Calvet, J., Computer-assisted analysis of the developing Purkinje neuron. I. Effects of the age of the animal at the moment of explantation on the subsequent dendritic development in organotypic cultures, 321
Calvet, M.-C. and Calvet, J., Computer-assisted analysis of the developing Purkinje neuron. II. A comparative study of the dendritic development in differently aged kittens and in organotypic cultures explanted at the same ages, 334
Casamenti, F., see Orzi, F., 99
Cavalheiro, E.A., see Ikonomidou-Turski, C., 275
Chalazonitis, A., see Chen, G.-G., 372
Chang, L.-R., see Tao, P.-L., 313
Chapman, C., see Zhao, B.-G., 62
Chen, G.-G., Chalazonitis, A., Shen, K.-F. and Crain, S.M., Inhibitor of cyclic AMP-dependent protein kinase blocks opioid-induced prolongation of the action potential of mouse sensory ganglion neurons in dissociated cell cultures, 372
Clarkson, T.W., see Aschner, M., 31
Conti, F., Fabri, M. and Manzoni, T., Immunocytochemical evidence for glutamatergic cortico-cortical connections in monkeys, 148
Coulter, D.A., see Disterhoft, J.F., 118
Crain, S.M., see Chen, G.-G., 372
Cridland, R.A. and Henry, J.L., Facilitation of the tail-flick reflex by noxious cutaneous stimulation in the rat: antagonism by a substance P analogue, 15
Cross, A.J., see Simpson, M.D.C., 76
Cupello, A., see Rapallino, M.V., 350

Deakin, J.F.W., see Simpson, M.D.C., 76
Dean, M.F., see Jenkins, H.G., 265
Deitmer, J.W. and Schlue, W.-R., Intracellular acidosis of identified leech neurones produced by substitution of external sodium, 233
Despres, G., see Romand, R., 167
Diamantstein, T., see Bindoni, M., 10
Diana, G., see Orzi, F., 99
Disterhoft, J.F., Golden, D.T., Read, H.R., Coulter, D.A. and Alkon, D.L., AHP reductions in rabbit hippocampal neurons during conditioning correlate with acquisition of the learned response, 118
Drachman, D.B., see Ramsay, D.A., 134
Dragunow, M., Murphy, K., Leslie, R.A. and Robertson, H.A., Localization of adenosine A_1-receptors to the terminals of the perforant path, 252
Drucker-Colin, R. and Prospéro-Carcía, O., Microinjection of carbachol into the pontine area is unable to modify insomnia induced by *p*-chlorophenylalanine (PCPA), 163
Duckrow, R.B., Regional cerebral flow and glucose utilization during hyperinsulinemia, 363

Enkvist, M.O.K., Holopainen, I. and Åkerman, K.E.O., The effect of K^+ and glutamate receptor agonists on the membrane potential of suspensions of primary cultures of rat astrocytes as measured with a cyanine dye, DiS-C2-(5), 67

Fabri, M., see Conti, F., 148
Farner, L.A., see Starcevic, V.P., 112
Feral, C., see Le Gall, S., 83
Ferrara, N., Ousley, F. and Gospodarowicz, D., Bovine brain astrocytes express basic fibroblast growth factor, a neurotropic and angiogenic mitogen, 223
Fieschi, C., see Orzi, F., 99
Freed, W.J., see Becker, J.B., 401
Freedman, R., see Parfitt, K.D., 242
Fukuda, A., see Minami, T., 258

Gariano, F.R. and Groves, P.M., Burst firing induced in midbrain dopamine neurons by stimulation of the medial prefrontal and anterior cingulate cortices, 194

Golden, D.T., see Disterhoft, J.F., 118
Gospodarowicz, D., see Ferrara, N., 223
Groves, P.M., see Gariano, F.R., 194

Hafidi, A., see Romand, R., 167
Harvey, A.R., Rush, R.A. and Keating, P.J., Cultured fetal tectal tissue grafted to the midbrain of newborn rats: morphology of grafts and innervation by host retinal and cortical axons, 89
Haseldonckx, M., see Van Reempts, J., 142
Heinemann, U., see Jones, R.S.G., 367
Henry, J.L., see Cridland, R.A., 15
Holopainen, I., see Enkvist, M.O.K., 67
Hosford, D.A. and McNamara, J.O., Microinjection of muscimol into entopeduncular nucleus suppresses pilocarpine but not maximal electroshock seizures in rats, 205
Hydén, H., see Rapallino, M.V., 350

Ikonomidou-Turski, C., Cavalheiro, E.A., Turski, L., Bortollotto, Z.A., Kleinrok, Z., Calderazzo-Filho, L.S. and Turski, W.A., Differential effects of non-steroidal anti-inflammatory drugs on seizures produced by pilocarpine in rats, 275

Jackson, I.M.D., see Amir, S., 160
Jacquin, M.F., see Renehan, W.E., 396
Jenkins, H.G., Martin, J. and Dean, M.F., Receptor-mediated uptake of β-glucuronidase into primary astrocytes and C6 glioma cells from rat brain, 265
Johnson,, R.F., Morin, L.P. and Moore, R.Y., Retinohypothalamic projections in the hamster and rat demonstrated using cholera toxin, 301
Johnston, G.A.R., see Lloyd, H.G.E., 391
Jones, R.S.G. and Heinemann, U., Verapamil blocks the afterhyperpolarization but not the spike frequency accommodation of rat CA1 pyramidal cells in vitro, 367
Jurson, P.A., see Robinson, T.E., 211
Jüch, P.J.W. and Rokx, J.T.M., Efferent projections of the parvocellular reticular nucleus to the mesencephalic trigeminal nucleus in rat, 185

Keating, P.J., see Harvey, A.R., 89
Keil, L.C., see Starcevic, V.P., 112
Keilhoff, G., see Šťastný, F., 56
Kleinrok, Z., see Ikonomidou-Turski, C., 275
Knowles, W.D., see Tesche, C.D., 190
Kokkoroyiannis, T., see Kuipers, U., 154
Krnjević, K., see Zhang, L., 174
Krusin-Elbaum, L., see Tesche, C.D., 190
Kuipers, U., Laouris, Y., Kokkoroyiannis, T., Meyer-Lohmann, J. and Windhorst, U., Relations between time- and frequency-domain measures of signal transmission from cutaneous afferents to dorsal horn neurons, 154

Landgraf, R., see Neumann, I., 181
Laouris, Y., see Kuipers, U., 154
Law, P.Y., see Tao, P.-L., 313
Le Gall, S., Feral, C., Van Minnen, J. and Marchand, C.R., Evidence for peptidergic innervation of the endocrine optic gland in Sepia by neurons showing FMRPamide-like immunoreactivity, 83
Leslie, R.A., see Dragunow, M., 252
Levitan, H., see Wadhawani, K.C., 22
Lisy, V., see Šťastný, F., 56
Litwak, M.S., see Arrezo, J.C., 286
Lloyd, H.G.E., Spence, I. and Johnston, G.A.R., Involvement of adenosine in synaptic depression induced by a brief period of hypoxia in isolated spinal cord of neonatal rat, 391
Loh, H.H., see Tao, P.-L., 313

Mann, D.M.A., see Simpson, M.D.C., 76
Manzoni, T., see Conti, F., 148
Marchand, C.R., see Le Gall, S., 83
Martin, J., see Jenkins, H.G., 265
Martínez-Murillo, R., see Villalba, R.M., 359
Mason, S.E., see Abraham, W.C., 40
McCall, R.D., see Renehan, W.E., 396
McNamara, J.O., see Hosford, D.A., 205
Mercuri, N.B., Calabresi, P. and Bernardi, G., Potassium ions play a role in the glycine-induced inhibition of rat substantia nigra zona compacta neurones, 199
Meyer-Lohmann, J., see Kuipers, U., 154
Minami, T., Oomura, Y., Nabekura, J. and Fukuda, A., Direct effects of 3,4-dihydroxybutanoic acid γ-lactone and 2,4,5-γ-trihydroxypentanoic acid γ-lactone on lateral and ventromedial hypothalamic neurons, 258
Moore, R.Y., see Johnson, R.F., 301
Morin, L.P., see Johnson, R.F., 301
Morrow, B.A., see Starcevic, V.P., 112
Murphy, K., see Dragunow, M., 252

Nabekura, J., see Minami, T., 258
Neumann, I., Schwarzberg, H. and Landgraf, R., Measurement of septal release of vasopressin and oxytocin by the push–pull technique following electrical stimulation of the paraventricular nucleus of rat, 181

Oomura, Y., see Minami, T., 258
Orzi, F., Diana, G., Casamenti, F., Palombo, E. and Fieschi, C., Local cerebral glucose utilization following unilateral and bilateral lesions of the nucleus basalis magnocellularis in the rat, 99
Ossowska, K., see Wardas, J., 378
Ousley, F., see Ferrara, N., 223

Palombo, E., see Orzi, F., 99
Parfitt, K.D., Freedman, R. and Bickford-Wimer, P.C., Electrophysiological effects of locally applied noradrenergic agents at cerebellar Purkinje neurons: receptor specificity, 242
Perciavalle, V., see Bindoni, M., 10
Pestronk, A., see Ramsay, D.A., 134
Prospéro-Garcia, O., see Drucker-Colin, R., 163
Ramsay, D.A., Drachman, D.B. and Pestronk, A., Rapid synthesis of acetylcholine receptors at neuromuscular junctions, 134
Rapallino, M.V., Cupello, A. and Hydén, H., Direct evidence for the presence of $GABA_A$ receptors on the cytoplasmic side of the Deiter's neurone membrane, 350
Rapoport, S.I., see Smith, Q.R., 1
Rapoport, S.I., see Wadhawani, K.C., 22
Read, H.R., see Disterhoft, J.F., 118
Renehan, W.E., Stansel, S.S., McCall, R.D., Rhoades, R.W. and Jacquin, M.F., An electron microscopic analysis of the morphology and connectivity of individual HRP-labeled slowly adapting vibrissa primary afferents in the adult rat, 396
Rhoades, R.W., see Renehan, W.E., 396
Robertson, D. and Winter, I.M., Cochlear nucleus inputs to

olivocochlear neurones revealed by combined anterograde and retrograde labelling in the guinea pig, 47
Robertson, H.A., see Dragunow, M., 252
Robinson, P.J., see Smith, Q.R., 1
Robinson, T.E., Jurson, P.A., Bennett, J.A. and Bentgen, K.M., Persistent sensitization of dopamine neurotransmission in ventral striatum (nucleus accumbens) produced by prior experience with (+)-amphetamine: a microdialysis study in freely moving rats, 211
Rodrigo, J., see Villalba, R.M., 359
Rokx, J.T.M., see Jüch, P.J.W., 185
Romand, R., Hafidi, A. and Despres, G., Immunocytochemical localization of neurofilament protein subunits in the spiral ganglion of the adult rat, 167
Rothe, F., see Šťastný, F., 56
Royston, M.C., see Simpson, M.D.C., 76
Rush, R.A., see Harvey, A.R., 89

Sawynok, J., see Sweeney, M., 346
Schlue, W.-R., see Deitmer, J.W., 233
Schmidt, W., see Šťastný, F., 56
Schroeder, C.E., see Arrezo, J.C., 286
Schwarzberg, H., see Neumann, I., 181
Severs, W.B., see Starcevic, V.P., 112
Shen, K.-F., see Chen, G.-G., 372
Simpson, M.D.C., Royston, M.C., Deakin, J.F.W., Cross, A.J., Mann, D.M.A. and Slater, P., Regional changes in [^{3}H]D-aspartate and [^{3}H]TCP binding site in in Alzheimer's disease brains, 76
Slater, P., see Simpson, M.D.C., 76
Smith, Q.R., Ziylan, Y.Z., Robinson, P.J. and Rapoport, S.I., Kinetics and distribution volumes for tracers of different sizes in the brain plasma space, 1
Soto, E. and Vega, R., Actions of excitatory amino acid agonists and antagonists on the primary afferents of the vestibular system of the axolotl (*Ambystoma mexicanum*), 104
Spence, I., see Lloyd, H.G.E., 391
Stansel, S.S., see Renehan, W.E., 396
Starcevic, V.P., Morrow, B.A., Farner, L.A., Keil, L.C. and Severs, W.B., Long-term recording of cerebrospinal fluid pressure in freely behaving rats, 112
Šťastný, F., Rothe, F., Schmidt, W., Wolf, G., Keilhoff, G. and Lisy, V., Changes in the activity of γ-glutamyl transpeptidase induced by kainic acid and surgical lesions of the hippocampal formation in young rats, 56
Sweeney, M., White, T. and Sawynok, J., 5-Hydroxytryptamine releases adenosine from primary afferent nerve terminals in the spinal cord, 346

Tao, P.-L., Chang, L.-R., Law, P.Y. and Loh, H.H., Decrease in δ-opioid receptor density in rat brain after chronic [D-Ala^2,D-Leu^5]enkephalin treatment, 313
Tesche, C.D., Krusin-Elbaum, L. and Knowles, W.D., Simultaneous measurement of magnetic and electric responses of in vitro hippocampal slices, 190
Thone, F., see Van Reempts, J., 142
Turski, L., see Ikonomidou-Turski, C., 275
Turski, W.A., see Ikonomidou-Turski, C., 275

Van Deuren, B., see Van Reempts, J., 142
Van Minnen, J., see Le Gall, S., 83
Van Reempts, J., Van Deuren, B., Haseldonckx, M., Van de Ven, M., Thone, F. and Bordergs, M., Purine nucleoside phosphorylase: a histochemical marker for glial cells, 142
Van de Ven, M., see Van Reempts, J., 142
Vega, R., see Soto, E., 104
Villalba, R.M., Martínez-Murillo, R., Blasco, I., Alvarez, F.J. and Rodrigo, J., C-PON containing neurons in the rat striatum are also positive for NADPH-diaphorase activity. A light microscopic study, 359

Wadhawani, K.C., Levitan, H. and Rapoport, S.I., Calcium transfer at endoneurial capillaries of the frog sciatic nerve, 22
Ward, D.G., Stimulation of the parabrachial nuclei with monosodium glutamate increases arterial pressure, 383
Wardas, J., Ossowska, K. and Wolfarth, S., Evidence for the independent role of GABA synapses of the zona incerta–lateral hypothalamic region in haloperidol-induced catalepsy, 378
West, T.E.G. and Wise, R.A., Effects of naltrexone on nucleus accumbens, lateral hypothalamic and ventral tegmental self-stimulation rate-frequency functions, 126
White, T., see Sweeney, M., 346
Windhorst, U., see Kuipers, U., 154
Winter, I.M., see Robertson, D., 47
Wise, R.A., see West, T.E.G., 126
Wolf, G., see Šťastný, F., 56
Wolfarth, S., see Wardas, J., 378

Zhang, L. and Krnjević, K., Intracellular injection of Ca^{2+} chelator does not affect spike repolarization of cat spinal motoneurons, 174
Zhao, B.-G., Chapman, C. and Bicknell, R.J., Functional κ-opioid receptors on oxytocin and vasopressin nerve terminals isolated from the rat neurohypophysis, 62
Ziylan, Y.Z., see Smith, Q.R., 1